

# Ambulatory N95 Frequency of Use Summary

N95s are only to be worn when caring for a confirmed COVID+/PUI patient or performing an aerosolizing procedure

High risk aerosolizing procedures	Low risk aerosolizing procedures	COVID+ patients & Non-aerosolizing procedures
Highly aerosolizing upper aerodigestive procedures	Procedures with lower risk for aerosol production	Episodic contact with COVID + patients
Place N95 in recollection bin upon completion of procedure	Keep your N95 in Tupperware container between uses. Place in recollection bin at the end of the shift	Only obtain an N95 when needed. Keep N95 in Tupperware between uses for 7 days after first use. After 7 days, place in recollection bin.
One N95 <u>per patient</u>	One N95 <u>per shift</u>	One N95 <u>per week</u>
<ul style="list-style-type: none"> <li>• Intubation</li> <li>• Extubation</li> <li>• ENT</li> <li>• Dental</li> <li>• BiPAP/CPAP</li> <li>• TEE</li> <li>• GI</li> </ul>	<ul style="list-style-type: none"> <li>• Nebulized medications</li> <li>• Suction</li> <li>• High flow nasal oxygen</li> <li>• Nasal swabs</li> </ul>	<ul style="list-style-type: none"> <li>• Exam room visits</li> <li>• Infusions</li> <li>• Radiology/Ultrasound Exams</li> <li>• Phlebotomy</li> </ul>