

MEDICATION ADHERENCE PACKAGING (MAP)

WHAT IS MEDICATION ADHERENCE PACKAGING?

- ❖ Pre-filled multi-dose medication packaging system
- ❖ Option to be organized by one, two, three or four times daily dosing
- ❖ Provides up to a 60 day supply of meds
Mail-order delivery preferred

PHARMACIES THAT OFFER MAP

- ❖ BMC Yawkey 850 Harrison Ave
Boston, MA
T: (617)-414-4883
- ❖ Cornerstone Health Solutions
41 Teed Drive Randolph, MA
T: (617)-414-4883

WHO MAY BENEFIT

- ❖ Patients on stable regimens (2 months without changes)
- ❖ Patients taking multiple medications
- ❖ Patients with difficulty organizing and understanding medications
- ❖ Patients who struggle with adherence

WHO MAY NOT BENEFIT

- ❖ Patients with unstable medication regimens (including PRN medications)
- ❖ Patients that live with small children (not childproof)
- ❖ Patients with unstable housing (mail order)
- ❖ Patients with unreliable form of communication with MAP team

OTHER CONSIDERATIONS

- ❖ Longer time to fill/receive compared to medications that are vialled
- ❖ Not available for discharge fills

- ❖ As needed and Schedule II medications are dispensed in vials
- ❖ Delivery to patient home by mail is preferred

HOW TO ENROLL

- ❖ Providers should speak with patients about the benefits and requirements of MAP and if it's right for them
- ❖ Patients should then be referred to the MAP team:
 - 617-414-0571
 - med.adherencepack@bmc.org
- ❖ Enrollment form will need to be completed

PATIENT COMMITMENT

- ❖ Patient must commit to having a monthly conversation with the MAP team to perform Medication Reconciliation prior to filling and shipping/picking-up order

PROVIDER COMMITMENT

- ❖ During each patient office visit, review medication list thoroughly in Epic and update medication list
- ❖ Contact pharmacy for all medication changes and direct patient to identify/remove/discard discontinued medications from MAP. Pharmacist available for patients requiring assistance with medication changes
- ❖ Consider delaying changed medication to next cycle. If immediately necessary, new dose will be dispensed in a vial until next cycle. Instruct patient to discard MAP dose

